

## Acta Tropica

Published by Elsevier

This article is being provided to filariasis.net with the permission of Acta Tropica and the Elsevier. You are free to download the article for education and teaching purposes, however, you must not post this article on any other web site or media, without first obtaining permission from the Editor of the Journal.

### About the Journal

*Acta Tropica*, is an international journal that covers biomedical and health sciences with particular emphasis on topics relevant to human and animal health in the tropics and the subtropics.

Its scope includes the biology of parasites and vectors, welcoming contributions concerning either basic or applied research in disciplines such as taxonomy, morphology, biochemistry, physiology and immunology; the development of tools for diagnosis and disease control; clinical and community medicine; and the epidemiology of communicable disease and health systems.

Contributions may be in the form of original papers, review articles or short communications - case reports are not considered for publication.

### Editors:

**A. Björkman, K. Berzins, F. Guhl**

See [editorial board](#) for all editors information

### Author enquiries:

Instructions to authors can be found in the first issue of every volume and online at <http://www.elsevier.com/locate/.actatropica>.

### Useful Links

#### Bibliographic & ordering Information

ISSN: 0001-706X

Imprint: ELSEVIER

Commenced publication 1944

Subscriptions for the year 2007, Volumes 101-104, 12 issues

#### Institutional online access:

[ScienceDirect eSelect](#)

For purchase of online access to this journal on ScienceDirect by credit card.

#### Institutional price:

[Order form](#)

JPY 216,200 for Japan

USD 1,829 for all countries

except Europe and Japan

EUR 1,637 for European countries

See also information about [conditions of sale & ordering procedures](#), and links to our [regional sales offices](#).

For an overview of recent dispatched issues, see the Journal issue [dispatch dates](#)

#### Audience

All clinicians and researchers dealing with tropical diseases, including parasitologists, microbiologists, immunologists and epidemiologists

#### Impact factor of this journal

2005: 1.80

- <sup>1</sup> East African Institute for Malaria and Other Vector-Borne Diseases, Amani, Tanzania  
<sup>2</sup> MRC Helminthiasis Research Unit, Tanga, Tanzania  
<sup>3</sup> Caribbean Epidemiology Centre, Port of Spain, Trinidad  
<sup>4</sup> Tropical Epidemiology Unit, London School of Hygiene and Tropical Medicine, U.K.

## Tanzania Filariasis Project

### Survey methodology and clinical manifestations of Bancroftian filariasis

P. WEGESA<sup>1</sup>, J. E. MCMAHON<sup>2</sup>, D. E. ABARU<sup>1</sup>, P. J. S. HAMILTON<sup>3</sup>,  
T. F. DE C. MARSHALL<sup>4</sup>, J. P. VAUGHAN<sup>4</sup>

Participating bodies: The East African Institute for Malaria and Other Vector-Borne Diseases, Amani, and the British Medical Research Council Helminthiasis Research Unit at Tanga. Other bodies associated and collaborating with the project since its commencement were: the World Health Organization, the Tanzania Ministry of Health, Department of Community Medicine (Medical Faculty, University of Dar es Salaam), London School of Hygiene and Tropical Medicine, and the Medical Research Centre at the Netherlands Royal Tropical Institute, Nairobi.

### Summary

In a Bancroftian filariasis survey on the coast of Tanzania microfilaria rates rose with age reaching 53% in the 60–69 year group followed by a slight fall above this age. The most important clinical manifestations were hydrocoele, funiculitis and elephantiasis, with hydrocoele presenting the most serious public health problem. Hydrocoele rates increased with age reaching 90% above the age of 70. The highest proportion of large hydrocoeles were also in the older age groups. A satisfactory methodology was established for the planning of future surveys leading to control.

*Key words:* Tanzania; Bancroftian filariasis; microfilaria; survey; hydrocoele.

### Introduction

Bancroftian filariasis surveys have verified high endemicity along the coasts of Tanzania (Jordan, 1956) and Kenya (Nelson et al., 1962) and in such

---

Correspondence: Dr. J. P. Vaughan, Department of Medical Statistics and Epidemiology, London School of Hygiene and Tropical Medicine, Keppel Street (Gower Street), London WC1E 7HT, U.K.




Fig. 1. The four divisions – Madanga, Pangani, Mwera and Mkwaja – surveyed in Pangani and the microfilaria (M), hydrocoele (H), and elephantiasis (E) rates in these divisions.

surveys, microfilaria (mf) prevalence rates have varied between 10 to 40% importance of hydrocoeles as a probable clinical manifestation of filariasis has been noted by Hawking (1940), Jordan (1955), Nelson et al. (1962), Men Kilama (1974), and Wijers and McMahan (1976).

In 1973 a Bancroftian filariasis project commenced in the Tanga region of Tanzania, which was aimed at a better understanding of the dynamic control of transmission.

The objectives of the present work were to gain information in the Pangani District of Tanga Region (Fig. 1) on:

1. the prevalence of mf in the general population;
2. the importance and frequency of the various clinical manifestations of filariasis; particular significance was given to investigating hydrocoeles as a health problem;

### 3. the testing of survey methodology and logistics.

The experience obtained in co-ordinating these clinico-parasitological surveys under field conditions was used to assess the degree of efficiency possible with the locally available administrative and technical services, and for the planning of future survey work.

## Methods and materials

### *Sampling and area surveyed*

Subjects from 12 randomly selected villages in 4 Divisions (Madanga, Pangani, Mwera and Mkwaja) in Pangani District (Fig. 1) were examined. Initially a 3 stage random cluster sampling method was used in which small samples of 50 subjects were examined. The stages were by selection of villages, then sections of villages and finally by Balozi<sup>5</sup>. This proved impractical and samples were increased to about 100 persons. In order to standardise with Wijers and Kinyanjui (1977) in Kenya those examined were males 10 years and over who (a) had been born in or came as a small baby to the village and who had never left the area except for occasional visits, or (b) claimed they had lived in the area for at least 10 years.

Because children were often absent from villages when examinations were conducted, in addition to a total of 787 males (6% of the total population in the Pangani District) examined in cluster samples, 504 pupils (males and females) aged 6–15 years from primary schools were also examined.

### *Parasitological*

In East Africa mf of *Wuchereria bancrofti* exhibit nocturnal periodicity. Because of problems associated with collecting blood at night and in order to standardise with a filariasis project being conducted in Kenya the provocative day test was used as described by Manson-Bahr and Wijers (1974) and in detail by McMahon et al. (1979). Briefly the method consisted of collecting 0.1 ml of finger prick blood in capillary tubes containing 3% acetic acid 50 min after oral administration of diethylcarbamazine base (DEC 100 mg to persons 15 years and above and 75 mg to those 10–14 years). Mf were counted in a counting chamber under the low power of a binocular microscope.

Fuglsang and Anderson (1974) found the number of mf of *Onchocerca volvulus* increased in the urine following diethylcarbamazine therapy. We investigated whether this occurred with *W. bancrofti* following the provocative test. Urine specimens were collected approximately 1 h after the DEC had been given and 10 ml samples were centrifuged and examined for mf.

### *Clinical*

Particular significance was placed on the histories of fever, lymphadenitis, lymphangitis, lymphadenopathy, chyluria and funiculitis as well as present clinical signs by means of a standardised questionnaire.

Hydrocoeles and groin glands were graded as follows:

*Hydrocoeles* – longitudinal axis: Grade I thickened cord or/and hydrocoele 6–8 cm, Grade II 8.1–11 cm, Grade III 11.1–15 cm, and Grade IV 15+ cm.

<sup>5</sup> In the political structure in villages in Tanzania, a senior person from a group of homes (called a Balozi or Ten House Chairman) is responsible for co-ordinating health and other activities of a small particular group. The Balozi was of great assistance in obtaining co-operation of the people selected for examination.

- Groin glands*<sup>6</sup> -I Not visible or diameter <3 cm  
II Visible or diameter > 3 cm  
III Skin fold present

The presence of elephantiasis (scrotal and limb) was recorded.

#### *Procedure*

In each village examinations were carried out in a suitable local home or administrative building. On the day prior to examination persons in the selected cluster sample were given cards which were presented at the examination. Blood samples were collected between 7.30–14.00 h.

The clinical examination was conducted by 2 medical officers (J.E.M. and D.E.). The importance of observer variation in clinical findings in onchocerciasis was emphasised by H et al. (1974). In the present work variation in clinical findings between the 2 clinicians was avoided by conducting duplicate clinical examination on the same subjects on randomly selected intervals throughout the survey. Subjects were re-examined in the event of any discrepancy. A close mutual agreement was reached by this method. At the end of the survey the rates and percentages recorded by each clinician for all lesions were compared.

The first microscopic examinations for mf were carried out at a field station in Pangar. Check examinations were later conducted on these specimens in the laboratories at Ta Amani.

#### *Period of survey*

The survey was conducted between June and August (1973). The seasonal variation in the period was not thought to be a problem since, in the case of Bancroftian filariasis, the long life span period of *W. bancrofti* plus the longevity of mf diminish the importance of the time between the commencement and termination of the survey.

## **Results**

The mf, hydrocoele and elephantiasis rates for the 4 divisions are shown in Fig. 1. As there were no marked differences in rates from different villages, for the further analysis the cluster samples have been combined.

#### *Microfilaria prevalence*

The mf prevalence rates by age for both the combined village cluster samples and schoolchildren are shown in Fig. 2. A rise of prevalence with age is evident in both groups followed by a slight fall in the oldest subjects in the village cluster samples.

#### *Fever*

There was no relation between microfilaraemia and fever either in persons having fever at the time of examination or in 108 persons with history of recent fever.

<sup>6</sup>As enlarged femoral and inguinal glands often coalesce they have been graded together as enlarged glands.




Fig. 2. Prevalence of microfilaraemia in village cluster samples and schools.

Table 1. Funiculitis grading for each age group

Age	No. examined	No. positive for funiculitis	%
10-14	74	7	9.5
15-19	75	24	36.9
20-29	160	49	30.6
30-39	146	45	30.8
40-49	102	39	38.2
50-59	89	34	38.2
60-69	106	41	38.7
70+	40	13	32.5
Totals	792	252	31.8

#### *Lymphatic system and hydrocoele*

Only 8 subjects (1%) had groin glands of Grade 3. Seven of these had a microfilaraemia.

Small glands grades 1 and 2 were found in 83% of subjects. Due to scabies, other skin infections, sores and wounds (many of the population do not wear shoes) lymphadenitis and lymphangitis were common. There were only 2 cases (0.2%) of retrograde lymphangitis. No cases of chyluria were seen.

A large proportion of postpubertal males gave a typical history of spermatic cord pain or had thickened and tender cords. The prevalence of funiculitis is shown in Table 1.




Fig. 3. Prevalence of hydrocoeles in village cluster samples.

The results shown in Fig. 3 indicate that the total hydrocoele prevalence rates increase with advancing age and Table 2 demonstrates that the proportion of larger hydrocoeles is greater in the older age groups.

Microfilaria prevalence in persons with normal testes and those with different grades of hydrocoele are shown in Table 3. There is no significant difference between these groups.

A similar pattern is shown in 33 subjects who had been operated upon for hydrocoeles, of whom 11 (33.3%) had microfilaraemias at the time of examination.

#### *Elephantiasis*

Prevalence rates of leg and scrotal elephantiasis were low, 2.8 and 1.5 respectively. The more severe lesions are commoner in older age groups. Elephantiasis was seen below the age of 19 years. No cases of arm elephantiasis were seen.

#### *Microfilaruria*

Only one case of microfilaruria was seen and this was in a child with

Table 2. Hydrocoele grading for each age group

Age	No. examined	Normal		Grade I		Grade II		Grade III		Grade IV	
		No.	%	No.	%	No.	%	No.	%	No.	%
10-14	76	72	94.7	3	3.9	1	1.3	-	-	-	-
15-19	66	52	78.8	10	15.1	4	6.0	-	-	-	-
20-29	160	122	76.3	23	14.4	15	9.4	-	-	-	-
30-39	147	88	59.9	31	21.1	20	13.6	8	5.4	-	-
40-49	102	35	34.3	30	29.4	32	31.4	3	3.0	2	2.0
50-59	89	29	32.6	22	24.7	31	34.8	5	5.6	2	2.2
60-69	106	25	23.6	28	26.4	37	35.0	13	12.3	3	2.8
70+	41	5	12.1	10	24.4	17	41.5	6	14.6	3	7.3
Totals	787	428	54.4	157	20.0	157	20.0	35	4.4	10	1.3

Table 3. Comparison of mf rates in subjects without (0) and with hydrocoeles Grades I-IV

	0	I	II	III	IV	Total I-IV
Total .....	428	157	157	35	10	359
No. with mf .....	164	66	63	13	3	145
Mf rate .....	38.3%	42.0%	40.1%	37.1%	30.0%	40.2%

maturia from concomitant urinary schistosomiasis. The provocative dose of diethylcarbamazine did not induce microfilaruria.

In all clinical manifestations there was a satisfactory correlation between findings of the 2 clinicians.

## Discussion

In the present work elephantiasis, funiculitis and hydrocoele were assumed to be the main clinical manifestations of Bancroftian filariasis. In comparison with the high hydrocoele rates, elephantiasis cannot be considered of major public health importance. In 1976 at Tanga hospital 15% of all major operations were for hydrocoele. Hydrocoele rates increased with age reaching nearly 90% in males over 70 years. Large hydrocoeles (Grades 3 and 4) occurred in 42 (9%) of subjects over 30 years, and resulted in 'dragging' and tiredness with a decreased ability to perform manual work. They are therefore an economic liability in a peasant community. Muhondwa (1978) noted that as well as having economic consequences hydrocoeles are considered to be a shameful condi-

tion and are socially unacceptable. Funiculitis may result in severe scrotal and the course of a typical attack was described by Wijers and McM (1976).

Fever and lymphadenopathy appeared to be of little value in this Bancroftian filariasis survey as the former is commonly due to malaria and skin infections commonly cause lymphadenopathy and lymphadenitis. Hanging and chyluria appear to be unimportant clinical complications of filariasis.

Important problems of methodology brought out in this survey were related to sampling, the provocative test, accurate identification of microfilariae, *W. bancrofti* and observer variation, both in clinical assessment and in recording of mf counts.

Prior to examining cluster samples of persons living in groups of houses it is important to have accurate census figures, since bias may result due to people with gross signs of filariasis, but not included in the cluster sample, wishing to be examined and attaching themselves to a household in the sample. Examination of the total population of villages in later surveys (McMahon, unpubl data) revealed that the age related rates for village cluster samples gave a good estimate of the mf and clinical prevalence patterns within the village.

The use of the daytime DEC provocative test was acceptable to subjects. A small number complained of pruritus of the skin. Because of the possibility of provoking Mazzotti reactions the test is contraindicated in onchocerciasis endemic areas. The test proved to be as sensitive in detecting microfilariae as the examination of night blood (McMahon et al., 1979). Its use in rural surveys is therefore justified. Although mf densities by day and night were highly correlated ( $r = 0.83$ ) they tended to be lower after provocative day DEC than in the corresponding night blood.

The mf of *W. bancrofti* are easily identifiable in the unstained cover slip chamber preparations. Apart from *W. bancrofti* the only human filarial species encountered was occasionally *Dipetalonema perstans*. In a few instances where small mf of *W. bancrofti* were difficult to differentiate from mf of *D. perstans* species identification was confirmed by making a stained film of finger prick blood from the subject concerned.

The regular but random cross checking of slides by technicians resulted in the maintenance of high standards of reading and the clinical cross check. Mutual discussions successfully established a high observer correlation between the 2 clinicians. Because of the good correlation established in clinical grading of hydrocoele and glands it was considered that the grading system was satisfactory for future surveys.

In conclusion, the main significance of Bancroftian filariasis in this area was shown to be the clinical manifestations of hydrocoele, funiculitis and lesser extent elephantiasis. A satisfactory methodology was established. Further research into the epidemiology of Bancroftian filariasis in this coastal area is required for the planning of future surveys leading to control.