

Screening of army soldiers for *Wuchereria bancrofti* infection in the metropolitan Recife region, Brazil: implications for epidemiological surveillance

Zulma Medeiros¹, Judilson Gomes², Fátima Bêliz¹, Amaury Coutinho¹, Patricia Dreyer³ and Gerusa Dreyer¹

¹ Departamento de Parasitologia do Centro de Pesquisas Aggeu Magalhães – Fundação Oswaldo Cruz, Recife, Brazil

² Seção do Serviço de Saúde da 7ª Região Militar e 7ª Divisão de Exército, Recife, Brazil

³ Universidade Federal de Pernambuco, Brazil.

Summary

Between 1989 and 1995, blood surveys were performed for *Wuchereria bancrofti* infection in several barracks of the Brazilian army in the metropolitan Recife region. For initial screening, 60 µl of capillary blood were examined for microfilaria. All men who tested positive had microfilaria quantified by filtration of venous blood through a polycarbonate membrane. Of 23773 men screened, 585 (2.5%) had microfilaria (mf). Microfilarial density ranged from < 1–8706 mf/ml of blood. Thirteen individuals had ultra-low microfilarial densities (1 mf/11 ml of blood). Characterization of 174 autochthonous cases made it possible to map 8 new districts in 4 cities within metropolitan Recife region where transmission of *W. bancrofti* was previously unknown. Routine screening of soldiers in the military may provide important surveillance data for national programmes to eliminate transmission of *W. bancrofti*.

keywords *Wuchereria bancrofti*; *Mansonella ozzardi*; public health; epidemiology; surveillance.

correspondence Dr Gerusa Dreyer, Departamento de Parasitologia – Centro de Pesquisas Aggeu Magalhães – FIOCRUZ, Av. Moraes Rego, s/No., Câmpus da UFPE, Cidade Universitária, Recife PE, CEP 50.670–420, Brazil.

Introduction

Although it has long been a neglected disease of the tropics, lymphatic filariasis has received increased attention in recent years with the development of improved diagnostic tools and the recognition that periodic single-dose mass treatment can profoundly suppress levels of microfilaria in the blood (Kessel *et al.* 1970; Mahoney & Kessel 1971; Eberhard *et al.* 1991; Andrade *et al.* 1995; Dreyer *et al.* 1995), thereby leading to interruption of transmission of the parasite (Panicker *et al.* 1991; Eberhard *et al.* 1992; Ismail *et al.* 1996; Kimura & Mataika 1996; Kazura *et al.* 1997; Bockarie *et al.* 1998). In May 1997 the World Health Assembly passed a resolution calling for the global elimination of lymphatic filariasis as a public health problem, and by November of that year, 13 countries had developed national filariasis elimination plans (WHO 1996; Ottesen *et al.* 1997).

One of the first major challenges facing these countries will be to determine the geographical distribution of filarial infection and to decide where filariasis elimination activities will be implemented. Surveillance for *Wuchereria bancrofti*

infection presents several potential difficulties. Unlike diseases for which elimination campaigns have been successful, such as smallpox, dracunculiasis and polio, most patent *W. bancrofti* infections are asymptomatic and many years may elapse between infection and development of obvious clinical sequelae such as hydrocele and lymphoedema. Thus, although rapid assessment techniques based on clinical disease have been proposed for lymphatic filariasis (Pani *et al.* 1995, 1997; Gyapong *et al.* 1996a, b, c, 1998), the sensitivity of these methods, if used alone, may be inadequate to detect infection in areas of low-level or recent transmission. Further, these techniques probably require training of many community-level health workers or deployment of mobile assessment teams. Particularly to identify foci of transmission in areas where lymphatic filariasis is not suspected to occur (making these areas likely to be overlooked or of low priority for rapid assessment), and for the purposes of certifying that *W. bancrofti* has been eliminated in a country, a routine, ongoing, 'passive' system is needed for detection of current *W. bancrofti* infection in a nationally representative sentinel population. Our experience in Recife, Brazil, where bancroft-

Z. Medeiros *et al.* Screening for *Wuchereria bancrofti* in the Brazilian army

ian filariasis is endemic (Dreyer 1987; Medeiros *et al.* 1992; Maciel *et al.* 1994, 1996), suggests that filarial screening of soldiers living or working in the military barracks may provide such a system.

Materials and methods

The project was approved by the Ethical Committee of Hospital das Clínicas – Universidade Federal de Pernambuco. Soldiers from 85 barracks in the metropolitan Recife region were targeted for screening. Before they were invited to have their blood examined for filarial infection, the soldiers attended an educational seminar that provided information on filarial infection, its diagnosis and prevention and signs and symptoms of acute and chronic filarial disease. After each soldier gave verbal consent to participate, information was collected on place of origin, current and past residential address and duration of residence in each location. Infections were considered autochthonous if they occurred in lifelong residents of metropolitan Recife region who had not travelled outside the area.

Fingerprick blood specimens were collected between 2300h and 0100h, the peak period of microfilaraemia in Metropolitan Recife Region (Dreyer *et al.* 1996a). For each soldier, three drops of capillary blood (approximately 60 µl) were placed on a glass slide, dried, stained with Carrazi haematoxylin-eosin, and examined under the microscope. To more accurately estimate microfilarial density in men who tested positive on the thick smear, one additional ml of peripheral blood was collected at night, filtered through a 3 µm polycarbonate membrane, and examined microscopically (Dennis & Kaen 1971). If microfilaria (mf) were not detected in 1 ml of filtered blood, this test was repeated by filtering 5 ml of blood, and if necessary, an additional 5 ml the following week. Every patient who tested positive for *W. bancrofti* mf was treated with diethylcarbamazine (DEC) or ivermectin by physicians working at the Military Hospital or the Filariasis Clinic of Centro de Pesquisas Aggeu Magalhães (CPqAM/FIOCRUZ), both in Recife.

Results

Between 1989 and 1995, 23773 men from 85 barracks in metropolitan Recife region (the cities of Recife, Jaboatão dos Guararapes and Olinda) were screened for microfilaraemia. None of the soldiers who attended the introductory lecture refused to undergo a blood examination. Mean age was 22 years (range: 18–58). 585 men (2.5%) were mf-positive (Table 1). Of these, 582 were infected by *W. bancrofti* while 3 individuals, aged 36, 48, and 39 years, were infected with *Mansonella ozzardi* (Manson 1897) between 1989 and 1991. These three men were lifelong residents of areas with known

Table 1 Prevalence of *W. bancrofti* and *M. ozzardi* infection among Army's military units in Metropolitan Recife Region, by year 1989–95

Year	N° of barracks	No. men examined	No. (%) with microfilaraemia†
1989	13	3594	147 (4.1)*
1990	07	2072	68 (3.3)
1991	16	3434	111 (3.2)**
1992	10	2346	60 (2.6)
1993	10	2664	57 (2.1)
1994	13	4489	74 (1.6)
1995	16	5174	68 (1.3)
Total	85	23773	585 (2.5)

† All infections are *W. bancrofti* except 3 cases represented by: *2 men infected by *M. ozzardi*; **1 man infected by *M. ozzardi*.

M. ozzardi transmission in Brazil (Rachou 1956). The annual filarial infection prevalence ranged from 1.3% to 4.1% (Table 1). The median interval between joining the army and screening was 6 months.

In 1989–91, the prevalence of *W. bancrofti* infection was significantly higher among soldiers 18–30-year-old (3.8%) than among those who were older (RR = 2.6, CI 1.5–4.6, $P = 0.00028$); 96% of infections were detected in soldiers aged 18–30 years (310/323). After the first year therefore screening was focused on military personnel < 30, with the exception of those who had been transferred from barracks elsewhere. Of 601 men who had been transferred from elsewhere and were examined between 1990 and 1991, 5 (0.8%), aged 31–58, had *W. bancrofti* microfilaraemia, 1 case being detected in 1990 and 4 cases in 1991 (Table 2).

Additional venous blood was collected from all 582 men with *W. bancrofti* detected in fingerprick samples and filtered for quantification of microfilarial density. All had *W. bancrofti* mf. The geometric mean microfilarial density was 239/ml (range: 1 mf/11 ml to 8,700/ml). 163 (28.0%) men had low-density microfilaraemia (1–100 per ml) and 13 (2.2%) ultra-low microfilarial densities (< 1 mf/ml). Geometric mean microfilarial density did not vary significantly between 1989 and 1995.

Of 582 men infected by *W. bancrofti*, only 174 (29.9%) had infections that were considered autochthonous to metropolitan Recife region (Figure 1). Other cases probably resulted from transmission there, but because the soldiers had lived in several areas within the metropolitan region, the precise location could not be determined. 165 soldiers with autochthonous infections resided in areas previously recognized to be endemic for filariasis (Jaboatão dos Guararapes, Olinda and Recife) and 9 lived in areas not previously recognized as having *W. bancrofti* transmission – Abreu e Lima (1

Table 2 Distribution of filarial infection among soldiers at military bases, by age

Age	mf + /all (%)								Total
	1989	1990	1991	1992	1993	1994	1995		
18-20	104/2175 (4.8)	59/1304 (4.5)	89/2047 (4.3)	53/1600 (3.3)	34/1450 (2.3)	45/2435 (1.8)	53/2664 (2.0)	437/13675 (3.2)	
21-30	33/1112 (3.0)	8/618 (1.3)	17/936 (1.8)	7/611 (1.1)	23/1006 (2.3)	29/1635 (1.8)	15/2068 (0.7)	132/7986 (1.6)	
31-40	5/197 (2.5)*	0/98	4/258 (1.5)*	0/91	0/121	0/253	0/277	9/1295 (0.7)	
41-50	5/103 (4.8)*	1/46 (2.2)	1/174 (0.6)	0/40	0/82	0/149	0/137	7/731 (1.0)	
>50	0/7	0/6	0/19	0/4	0/5	0/17	0/28	0/86	
Total	147/3594 (4.1)	68/2072 (3.3)	111/3434 (3.2)	60/2346 (2.6)	57/2664 (2.1)	74/4489 (1.6)	68/5174 (1.3)	585/23773 (2.5)	

* Including cases of *M. ozzardi*.

case from Caetes district), Cabo de Santo Agostinho (1 case from Ponte dos Carvalhos), Camaragibe (2 cases from Fabrica district, 1 case from Bairro dos Estados district and 1 case from Bairro Novo district), and Paulista (1 case from Maranguape district, 1 case from Janga district and 1 case from Paratibe district). Subsequent follow-up of these cases

and their family members documented transmission in these areas (Zulma Medeiros, unpublished data).

Discussion

Surveillance systems for lymphatic filariasis are urgently




Figure 1 Distribution of autochthonous cases of lymphatic filariasis in the region of metropolitan Recife, northeastern Brazil. □ Previously recognised foci; ▨ Newly discovered foci.

needed to support and guide the global elimination programme recently announced by the World Health Organization (Ottesen & Ramachandran 1995; WHO 1995; Ottesen *et al.* 1997). It is likely that no single type of surveillance will meet the needs of all filariasis-endemic countries or of filariasis elimination programmes in all phases of their evolution (e.g. initial geographical assessment, programme monitoring, and certification of elimination). Recent work to develop techniques for initial filariasis assessment has focused on the use of clinical signs, particularly hydrocele, as indirect indicators of transmission (Pani *et al.* 1995, 1997; Gyapong *et al.* 1996a, b, c, 1998), or on collection and testing of fingerprick blood specimens for *W. bancrofti* infection by mobile survey teams. Our work with soldiers in the military as part of the lymphatic filariasis research and treatment programme at CPqAM in Recife, Brazil suggests that routine annual screening of military recruits is feasible and may complement these other surveillance methods. This type of ongoing routine surveillance may play a useful role in identifying new areas of transmission and in providing a back-up system that can enhance the level of certainty among public health officials when it appears that filariasis is being eliminated.

The military offers a convenient and fairly representative population for surveillance and assessment of a variety of diseases (Ali *et al.* 1985; Gill & Bailey 1989; Sharp *et al.* 1995; Drabick *et al.* 1997; Eitzen & Sawyer 1997). For example, serosurveys of military populations in the United States have been used to determine exposure (seropositivity) to *Toxoplasma gondii* (Smith *et al.* 1996). Several features of military bases also make them attractive for the purposes of screening and epidemiological surveillance for *W. bancrofti* infection. First, young adult males represent an ideal screening population because of their elevated risk for *W. bancrofti* infection and microfilaraemia. In most populations where lymphatic filariasis is endemic, the prevalence of microfilaraemia tends to increase with age to about 20 years, after which it remains consistently high (Brabin 1990; Lammie *et al.* 1994; Albuquerque *et al.* 1995; Chanteau *et al.* 1995). In addition, during the reproductive years, the prevalence of microfilaraemia tends to be higher in males than in females (Brabin 1990; Pani *et al.* 1991; Albuquerque *et al.* 1995). The prevalence of chronic filarial disease, such as hydrocele, tends to increase with age; thus most infected young males are asymptomatic (Ottesen 1993; Norões *et al.* 1996). Unless screening is offered to these men, their infections go undetected and they remain untreated. Without effective treatment, they serve as reservoirs for further transmission and their lymphatic damage progresses, thereby increasing their risk of clinical disease.

Secondly, since lymphatic filariasis causes hydrocele, it is a disease of considerable concern to the military authorities and to the soldiers themselves. In Brazil and in other

countries, hydrocele is one of several criteria for exclusion from military service. Soldiers often expressed keen interest in lymphatic filariasis; after they were tested at the military base, for example, many of them brought their entire families to the Filariasis Clinic at CPqAM/FIOCRUZ to be screened for *W. bancrofti* infection.

Third, screening military recruits is convenient, safe, and logistically uncomplicated. In Brazil, prospective soldiers are required to undergo a physical examination and screening for certain diseases before being allowed to enter the army. Currently, no testing for microfilaraemia is required, but it would be a simple matter to collect fingerprick blood for detection of mf. In our surveys, soldiers were highly cooperative with this procedure, and medical staff at the base were eager to learn how to diagnose lymphatic filariasis. In fact, in 1992, army personnel began enrolling in courses at CPqAM/FIOCRUZ, which qualified them to diagnose and treat *W. bancrofti* microfilaraemia cases. This was the first time a programme for diagnosis and treatment of an endemic disease was adopted by the Army in Brazil. Nighttime collection of blood on the military base was easy to arrange, and, unlike nighttime blood collection in some peri-urban slums, where lymphatic filariasis is emerging as an important public health problem, it was also safe for the blood collection team. Further, collecting blood at the peak time of microfilarial density, between 2300h and 0100h, was much more feasible on the army base than it is in most communities. Thus, the sensitivity of screening was enhanced.

Finally, at least in Brazil, military recruits are considered a representative cross-section of the young male population in the region where the base is located. Soldiers tend to be stationed at bases located near their town of residence. Screening of army soldiers has already been used elsewhere in Brazil to successfully identify foci of transmission of *W. bancrofti* in areas that were not considered to be endemic. For example, in 1990, two autochthonous *W. bancrofti* infections were detected when soldiers were screened at the 59^o Batalhão de Infantaria Motorizada in Maceió-Alagoas-Brazil (Dreyer *et al.* 1991). At the time, Maceió was considered by the Ministry of Health not to be endemic for *W. bancrofti* (Ministério da Saúde Brasil 1986). As a result of these findings, however, an epidemiological investigation was begun, active transmission was confirmed, and the Health Ministry recognized that filariasis was indeed a public health problem in Maceió. Filariasis research and control programmes have already begun in this area (Fontes *et al.* 1994, 1998).

Five years ago, *W. bancrofti* was also considered to be under control in Recife (Ministério da Saúde Brasil 1986). Work by Dreyer and colleagues subsequently documented transmission in 1985 (Dreyer 1987; Medeiros *et al.* 1992; Dreyer *et al.* 1995; 1996). Lymphatic filariasis is now considered an important public health problem of metropolitan

Recife region, with an overall prevalence of ~1%, but with prevalences of up to 15% in some districts (Maciel *et al.* 1996). Because active case detection and surveillance for *W. bancrofti* infection have only begun in recent years, it is difficult to say with absolute certainty whether transmission of this parasite is increasing in intensity and spreading to new areas, but this appears to be the case.

Our screening of military recruits identified 8 districts within four new cities in metropolitan Recife with active transmission of *W. bancrofti*. Autochthonous cases of *W. bancrofti* infection were identified in lifelong residents of a total of 7 cities in metropolitan Recife region; these infected persons resided in 79 districts of the metropolitan area (Figure 1).

Recently, considerable attention has focused on new circulating antigen-detection assays for diagnosis of *W. bancrofti* infection (More & Copeman 1990; Rocha *et al.* 1996; Freedman *et al.* 1997; Nicolas 1997; Weil *et al.* 1997). These assays have a high sensitivity and specificity for bancroftian filariasis infection. However, because the antigen is primarily adult worm-related, caution is warranted in using them for monitoring control programmes where the objective is to reduce or interrupt transmission. Transmission is directly associated with microfilaraemia, and for the first few years after treatment with antifilarial drugs, marked reductions in microfilaraemia are observed, but the prevalence of antigenemia remains constant (Weil *et al.* 1988). Therefore, monitoring mf levels is likely the best option for assessing the effectiveness of treatment in reducing transmission, particularly in the short term. Another drawback of the antigen assays is the fact that they are relatively expensive and their purchase on the international market requires 'hard currency'.

Although antigen-and even PCR-based assays have many advantages and are likely to be used more frequently in the diagnosis of lymphatic filariasis, detection of microfilaraemia using fingerprick blood will probably continue to play an important role in epidemiological surveillance and in monitoring the effectiveness of interventions to interrupt transmission (Nicolas 1997; Weil *et al.* 1997). Although we were unable to measure the sensitivity of the thick smear in this screening programme, it appeared to be relatively high, in view of the fact that 13 of men who had microfilaraemia detected in fingerprick samples had ultra-low microfilarial densities (< 1/ml venous blood). In part, this may have been because we examined 3 drops (60 µl), rather than a single drop (20 µl) of capillary blood. Previous work showed that, among persons with < 100 mf/ml, the sensitivity of thick smear increases from 68% to 94% as the volume of examined blood increases from 20 µl to 60 µl (Dreyer *et al.* 1996). Therefore we recommend, not only for the military but also for the national filariasis elimination programme of Health

Ministry (Fundação Nacional de Saúde –FNS), that 3 drops (60 µl) of capillary blood be collected by fingerprick for examination of microfilaraemia.

In summary, through routine screening during a 7-year period, we provided health education to thousands of soldiers, identified and treated 582 men 'silently' infected with *W. bancrofti*, and detected several new areas of transmission in metropolitan Recife. Our experience with the army suggests that the military services, including the Navy, Air Force and Military Police, may provide an excellent opportunity for ongoing routine surveillance of *W. bancrofti* infection at the national level. Such a surveillance system can be used by national filariasis elimination programmes to identify new areas of transmission and to monitor trends in prevalence over time in areas where transmission is known to occur. To be useful, military-based surveillance for *W. bancrofti* infection should be linked to thorough and timely case follow-up, particularly when infections are detected in residents of areas where transmission is currently not known to occur. This kind of surveillance should not replace community- or district-level initial assessment and monitoring of interventions to interrupt transmission. However, military-based surveillance can provide an adjunct surveillance system at the national level, particularly for the purpose of certifying countries as free of *W. bancrofti* transmission.

Acknowledgements

The authors are indebted to the General Commander of 7^a Região Militar e da 7^a Divisão de Exército for making this work possible in the period of 1989–97; to commanders of the barracks; health workers from the barracks and from Hospital Geral do Recife; to the soldiers who were examined and demonstrated enthusiasm during the accomplishment of the entire work; to Dr David Addiss for the critical reading and helpful discussion of the manuscript. This study was partially financed by UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Disease (ID number 880533 E30/181/16), by FACEPE (ID number 0157.2.13/94), by FIOCRUZ (PAPES n^o 2, PAP number 01938–8) and by OPAS (ID number ASC 97/00181–0)

References

- Albuquerque MFM, Marzochi MC, Ximenes RA, Braga MC, Silva MCM & Furtado AF (1995) Bancroftian filariasis in two urban areas of Recife, Brazil: The role of individual risk factors. *Revista do Instituto de Medicina tropical de São Paulo* 37, 225–233.
- Ali EM, Abdel-Wahab F & Romia SA (1985) A preliminary survey of bancroftian filariasis among soldiers in Mansoura, Egypt. *Journal of the Egyptian Society of Parasitology* 15, 493–498.

Z. Medeiros *et al.* **Screening for *Wuchereria bancrofti* in the Brazilian army**

- Andrade LD, Medeiros Z, Pires ML *et al.* (1995) Comparative efficacy of three different diethylcarbamazine regimens in lymphatic filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **89**, 319–321.
- Bockarie M, Alexander N, Hyun P *et al.* (1998) Randomised community-based trial of annual single-dose diethylcarbamazine with or without ivermectin against *Wuchereria bancrofti* infection in human beings and mosquitoes. *The Lancet* **351**, 162–168.
- Brabin L (1990) Sex differentials in susceptibility to lymphatic filariasis and implications for maternal child immunity. *Epidemiology and Infection* **105**, 335–353.
- Chanteau S, Glaziou P, Plichart C *et al.* (1995) *Wuchereria bancrofti* filariasis in French Polynesia: age-specific patterns of microfilaremia, circulation antigen, and specific IgG and IgG4 responses according to transmission level. *International Journal of Parasitology* **25**, 81–85.
- Dennis DT & Kaen BH (1971) Isolation of microfilariae: report of a new method. *Journal of Parasitology* **57**, 1146–1147.
- Drabick JJ, Gambel JM, Gouvea VS *et al.* (1997) A cluster of acute hepatitis E infection in United Nations Bangladeshi peacekeepers in Haiti. *American Journal of Tropical Medicine and Hygiene* **57**, 449–454.
- Dreyer G (1987) Filariasis Programme. *Memórias Do Instituto Oswaldo Cruz* **82**, 359–360.
- Dreyer G, Coutinho A, Miranda D *et al.* (1995) Treatment of bancroftian filariasis in Recife, Brazil: a two year comparative study of the efficacy of single treatments with ivermectin or diethylcarbamazine. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **89**, 98–102.
- Dreyer G, Medeiros Z, Bêliz F *et al.* (1991) Autochthonous *Wuchereria bancrofti* microfilaraemia in the city of Maceió, Alagoas – Brazil. *Memórias Do Instituto Oswaldo Cruz* **86**, 495–496.
- Dreyer G, Pimentel A, Medeiros Z *et al.* (1996a) Studies on the periodicity and intravascular distribution of *Wuchereria bancrofti* microfilariae in paired samples of capillary and venous blood from Recife, Brazil. *Tropical Medicine and International Health* **1**, 264–272.
- Eberhard ML, Dickerson JW, Hightower AW & Lammie PJ (1991) Bancroftian filariasis: long-term effects of treatment with diethylcarbamazine in a Haitian population. *American Journal of Tropical Medicine and Hygiene* **45**, 728–733.
- Eberhard ML, Hightower AW, McNeeley DF & Lammie PJ (1992) Long-term suppression of microfilaraemia following ivermectin treatment. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **86**, 287–288.
- Eitzen JP & Sawyer RG (1997) Sexually transmitted diseases: risk behaviors of female active duty US army recruits. *Military Medicine* **162**, 686–689.
- Fontes G, Brito AC, Calheiro CML, Antunes CM & Rocha EMM (1994) Situação atual da filariose bancroftiana na cidade de Maceió, Estado de Alagoas, Brasil. *Cadernos de Saúde Pública* **10**, 293–300.
- Fontes G, Rocha EMM, Brito AC & Antunes CMF (1998) Lymphatic filariasis in Brazilian urban area (Maceió, Alagoas) *Memórias do Instituto Oswaldo Cruz* **96**, 705–710.
- Freedman DO, De Almeida A, Miranda J, Plier AD & Braga C (1997) Field trial of a rapid card test for *Wuchereria bancrofti*. *Lancet* **350**, 1681.
- Gill GV & Bailey JW (1989) Eosinophilia as a marker for chronic strongyloidiasis – use of a serum ELISA test to detect asymptomatic cases. *Annals of Tropical Medicine and Parasitology* **83**, 249–252.
- Gyapong JO, Adjei S & Sackey SO (1996a) Descriptive epidemiology of lymphatic filariasis in Ghana. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **90**, 26–30.
- Gyapong JO, Adjei S, Gyapong M & Asamoah G (1996b) Rapid community diagnosis of lymphatic filariasis. *Acta Tropica* **61**, 65–74.
- Gyapong JO, Gyapong M & Adjei S (1996c) The epidemiology of acute adenolymphangitis due to lymphatic filariasis in Northern Guyana. *American Journal of Tropical Medicine and Hygiene* **54**, 591–595.
- Gyapong JO, Webber RH, Morris JO & Bennett S (1998) Prevalence of hydrocele as a rapid diagnostic index for lymphatic filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **92**, 40–43.
- Ismail MM, Weil GJ, Jayasinghe KSA *et al.* (1996) Prolonged clearance of microfilaraemia in patients with bancroftian filariasis after multiple high doses of ivermectin or diethylcarbamazine. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **90**, 684–688.
- Kazura J, Bockarie M, Alexander N *et al.* (1997) Transmission intensity and its relationship to infection and disease due to *Wuchereria bancrofti* in Papua New Guinea. *Journal of Infectious Diseases* **176**, 242–246.
- Kessel JF, Silica N, Tompkins H & Jones K (1970) Periodic mass treatment with diethylcarbamazine for the control of filariasis in American Samoa. *Bulletin of the World Health Organization* **43**, 817–825.
- Kimura E & Mataika JU (1996) Control of lymphatic filariasis by annual single-dose diethylcarbamazine treatments. *Parasitology Today* **12**, 240–244.
- Lammie PJ, Hightower AW & Eberhard ML (1994) Age-specific prevalence of antigenemia in a *Wuchereria bancrofti* – exposed population. *American Journal of Tropical Medicine and Hygiene* **51**, 348–355.
- Maciel MAV, Marzochi KBF, Silva EC, Rocha A & Furtado AF (1994) Estudo comparativo de áreas endêmicas de filariose bancroftiana na RMR, Brasil. *Cadernos de Saúde Pública* **10** (Suppl.), 301–309.
- Maciel A, Rocha A, Marzochi KB *et al.* (1996) Epidemiological study of bancroftian filariasis in Recife. *Northeastern Brazil. Memórias do Instituto Oswaldo Cruz* **91**, 449–455.
- Mahoney LE & Kessel JT (1971) Treatment failure in filariasis mass treatment programmes. *Bulletin of the World Health Organization* **45**, 35–42.
- Medeiros Z, Dreyer G, Andrade L, Pires ML, Mendes J & Pimentel R (1992) *Wuchereria bancrofti* microfilarial density of autochthonous cases and natural *Culex* infectivity rates in Northeast Brazil. *Journal of Tropical Medicine and Hygiene* **95**, 214–217.
- MS – Ministério da Saúde – Brasil (1986) Superintendência de Campanhas de Saúde Pública. Departamento de Erradicação e

Z. Medeiros *et al.* **Screening for *Wuchereria bancrofti* in the Brazilian army**

- Controle de Endemias. Demonstrativo dos Resultados obtidos em 1985 e projeções para 1986. Brasília – DF, pp. 83–87.
- More SJ & Copeman DB (1990) A highly specific and sensitive monoclonal antibody-based ELISA for the detection of circulating antigen in bancroftian filariasis. *Tropical Medicine and Parasitology* **41**, 403–406.
- Nicolas L (1997) New tools for diagnosis and monitoring of bancroftian filariasis parasitism: the Polynesian experience. *Parasitology Today* **13**, 370–375.
- Norões J, Addiss D, Santos A, Medeiros Z, Coutinho A & Dreyer G (1996) Ultrasonographic evidence of abnormal lymphatic vessels in young men with adult *Wuchereria bancrofti* infection in the scrotal area. *The Journal of Urology* **156**, 409–412.
- Ottesen EA (1993) Filarial infections. *Infectious Disease Clinics of North America* **7**, 619–633.
- Ottesen EA & Ramachandran CP (1995) Lymphatic filariasis infection and disease: control strategies. *Parasitology Today* **11**, 129–131.
- Ottesen EA, Duke BOL, Karan M & Behbehani K (1997) Strategies and tools for the control/elimination of lymphatic filariasis. *Bulletin of World Health Organization* **75**, 491–503.
- Pani SP, Balakrishnan N, Srividya A, Bundy DA & Grenfell BT (1991) Clinical epidemiology of bancroftian filariasis: effect of age and gender. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **85**, 260–264.
- Pani SP, Yuvaraj J, Vanamail P *et al.* (1995) Episodic adenolymphangitis and lymphedema in patients with bancroftian filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **89**, 72–74.
- Pani SP, Srividya A, Krishnamoorthy K, Das PK & Dhanda V (1997) Rapid assessment procedures (RAP) for lymphatic filariasis. *National Medical Journal of India* **10**, 19–22.
- Panicker KN, Krishnamoorthy K, Sabesan S, Prathida J & Abidha J (1991) Comparison of effects of mass annual and biannual single dose therapy with diethylcarbamazine for the control of malayan filariasis. *Southeast Asian Journal of Tropical Medicine and Public Health* **22**, 402–411.
- Rachou RG (1956) Distribuição geográfica das filárias humanas no Brasil. *Revista Brasileira de Malariologia e Doenças Tropicais* **9**, 79–100.
- Rocha A, Addiss D, Ribeiro ME *et al.* (1996) Evaluation of the Og4C3 ELISA in *Wuchereria bancrofti* infection: Infected persons with undetectable or ultra-low microfilarial densities. *Tropical Medicine and International Health* **1**, 859–864.
- Sharp TW, Wallace MR, Hayes CG *et al.* (1995) Dengue fever in US troops during operation Restore Hope, Somalia. 1992–93. *American Journal of Tropical Medicine and Hygiene* **53**, 89–94.
- Smith KL, Wilson M, Hightower AW *et al.* (1996) Prevalence of *Toxoplasma gondii* antibodies in US military recruits in 1989: comparison with data published in 1965. *Clinical Infectious Diseases* **23**, 1182–1183.
- Weil GJ, Lammie PJ & Weiss N (1997) The ICT filariasis test: a rapid-format antigen test for diagnosis of bancroftian filariasis. *Parasitology Today* **13**, 401–404.
- Weil GJ, Sethumadhavan KVP, Santhanam S, Jain DC & Ghosh TK (1988) Persistence of parasite antigenemia following diethylcarbamazine therapy of bancroftian filariasis. *American Journal of Tropical Medicine and Hygiene* **38**, 589–595.
- World Health Organization (1995) Lymphatic filariasis: an eradicable disease. *World Health Organization Drug Information* **9**, 135–138.
- World Health Organization (1996) TDR NEWS – UNDP/WORLD BANK/WHO Special Programme for Research and Training in Tropical Disease, No 49 – March.